


This article was downloaded by: [VPI Virginia Polytechnic Institute & State University]

On: 23 May 2011

Access details: Access Details: [subscription number 769791923]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Natural Product Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713398545>

### Antioxidant activity and total phenolic content of aqueous and methanolic extracts of Jordanian plants: an ICBG project

Feras Q. Alali<sup>ab</sup>; Khaled Tawaha<sup>c</sup>; Tamam El-Elimat<sup>b</sup>; Maha Syouf<sup>d</sup>; Mosa El-Fayad<sup>d</sup>; Khaled Abulaila<sup>d</sup>; Samara Joy Nielsen<sup>e</sup>; William D. Wheaton<sup>f</sup>; Joseph O. Falkinham III<sup>f</sup>; Nicholas H. Oberlies<sup>g</sup>

<sup>a</sup> Faculty of Pharmacy, Department of Medicinal Chemistry and Pharmacognosy, Jordan University of Science and Technology, Irbid 22110, Jordan <sup>b</sup> Pharmaceutical Research Center, Jordan University of Science and Technology, Irbid 22110, Jordan <sup>c</sup> Faculty of Pharmacy, Department of Pharmaceutical Sciences, University of Jordan, Amman, Jordan <sup>d</sup> National Center for Agricultural Research and Technology Transfer, Baqa' 19381, Jordan <sup>e</sup> Social and Statistical Sciences, Research Triangle Institute, North Carolina, USA <sup>f</sup> Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA <sup>g</sup> Natural Products Laboratory, Research Triangle Institute, North Carolina, USA

**To cite this Article** Alali, Feras Q. , Tawaha, Khaled , El-Elimat, Tamam , Syouf, Maha , El-Fayad, Mosa , Abulaila, Khaled , Nielsen, Samara Joy , Wheaton, William D. , Falkinham III, Joseph O. and Oberlies, Nicholas H.(2007) 'Antioxidant activity and total phenolic content of aqueous and methanolic extracts of Jordanian plants: an ICBG project', *Natural Product Research*, 21: 12, 1121 – 1131

**To link to this Article:** DOI: 10.1080/14786410701590285

**URL:** <http://dx.doi.org/10.1080/14786410701590285>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Antioxidant activity and total phenolic content of aqueous and methanolic extracts of Jordanian plants: an ICBG project

FERAS Q. ALALI\*†‡, KHALED TAWAHA§, TAMAM EL-ELIMAT‡,  
MAHA SYOUF¶, MOSA EL-FAYAD¶, KHALED ABULAILA¶,  
SAMARA JOY NIELSEN⊥, WILLIAM D. WHEATON§,  
JOSEPH O. FALKINHAM III§ and NICHOLAS H. OBERLIES¥

†Faculty of Pharmacy, Department of Medicinal Chemistry and Pharmacognosy,  
Jordan University of Science and Technology, P.O. Box 3030, Irbid 22110, Jordan

‡Pharmaceutical Research Center, Jordan University of Science and Technology,  
P.O. Box 3030, Irbid 22110, Jordan

§Faculty of Pharmacy, Department of Pharmaceutical Sciences, University of Jordan,  
Amman 11942, Jordan

¶National Center for Agricultural Research and Technology Transfer,  
P.O. Box 639, Baqa' 19381, Jordan

||Geospatial Science and Technology Program Research Triangle Institute, P.O.Box 12194,  
Research Triangle Park, North Carolina 27709, USA

⊥Social and Statistical Sciences, Research Triangle Institute, P.O.Box 12194, Research  
Triangle Park, North Carolina 27709, USA

§Department of Biological Sciences, Virginia Polytechnic Institute and State University,  
Blacksburg, Virginia 24061, USA

¥Natural Products Laboratory, Research Triangle Institute, P.O. Box 12194, Research  
Triangle Park, North Carolina 27709, USA

(Received 14 February 2007; in final form 19 June 2007)

As part of an International Cooperative Biodiversity Groups (ICBG) program to study Jordan's biodiversity, the relative levels of antioxidant activity and the total phenolic content of aqueous and methanolic extracts of a total of 95 plant species, all of Jordanian origin and those collected at random, have been measured. The total phenolic content of aqueous and methanolic extracts of the investigated plant species ranged from 4.4 to 78.3 mg and from 2.1 to 52.8 mg gallic acid equivalents  $g^{-1}$  dry weight, respectively, while the total antioxidant capacity ranged from 20.0 to 916.7 and from 15.1 to 915.6  $\mu$ mol Trolox equivalents  $g^{-1}$  dry weight, respectively. Based on this collection, approximately 5% of assayed plants showed high levels of antioxidant activity. There was a significant linear correlation between antioxidant activity and total phenolic content for aqueous and methanolic extracts, suggesting that phenolic compounds were the predominant antioxidant components in the investigated plant species. Interestingly, a few of the collected plants had high-antioxidant activity yet "low" phenolic content includes *Ceratonia siliqua* and *Viscum cruciatum*. These plants may serve as sources of antioxidants with new chemotypes.

**Keywords:** Antioxidants; Total phenolic content; Jordanian flora; Free radicals

\*Corresponding author. Tel.: +962-2-7201000. Fax: +962-2-7201075. Email: falali@just.edu.jo

## 1. Introduction

The Hashemite Kingdom of Jordan (Jordan) lies in the heart of the Middle East, about 100 km from the southeastern coast of the Mediterranean Sea, between latitudes 29°–33°N and longitudes 35°–39°E. It covers a wide range in elevation, from the lowest place on earth, 400 m below sea level near the Dead Sea, to plateaus of more than 1700 m above sea level near the Jebel Rum. This dynamic topography and climate places the geopolitical borders of Jordan at the junction of four biogeographical areas: the Mediterranean, the Irano-Turanian, the Saharo-Arabian, and the Tropical or Sudanian (figure 1) [1,2]. As such, Jordan encompasses a unique and rich habitat for a wide variety of plant life, most of which can be collected and studied efficiently in a relatively small land area [3,4]. More than 2500 wild plant species from 700 genera are found in Jordan, and of these, there are approximately 100 endemic species, 250 rare species, and 125 very rare species [5].

In 2003, Research Triangle Institute (RTI) in collaboration with Jordan University of Science and Technology, and Virginia Polytechnic Institute and State University, received a planning grant from the Fogarty International Center of the US National Institutes of Health, as part of the International Cooperative Biodiversity Groups (ICBG) program, to promote sustainable economic growth, biodiversity conservation, and training and education in Jordan through natural products-based drug discovery. The team focused on plants and nonobligate predatory soil bacteria as potential sources for new drug leads, in particular, to compounds, acting as anticancer agents,




Figure 1. Jordan's biogeographic regions and plant collection sites.

antibiotics, and/or modulators of the central nervous system. These studies are unique, in that very few plants of Middle Eastern origin, particularly those of Jordan, have been explored for pharmaceutical leads, especially from the extreme environments found in the deserts and near the Dead Sea. Since 80% of plants found in Jordan are common with other countries of the Middle East, Jordan's flora may be considered as the representative of the larger region [5].

The team collected over 120 plant samples, roughly 25–30 from each of the four biogeographic regions of Jordan (figure 1). At this early stage in the program, random sampling was chosen (as opposed to targeted collections) to demonstrate proof of concept that Jordan was a viable source of plants with promising biological activities. The sampling strategy included a linear transect of Jordan, focusing on the western portion of Jordan near the Jordan Valley, to assay the biodiversity and pharmaceutical potential of all four biogeographic regions. Several of the collected samples displayed promising biological activities, some of which have been reported previously [6–8] and some of which are currently under investigation. Herein, we present the antioxidant activity and total phenolic content for 95 plant samples, and statistical correlations are drawn between such activities for both aqueous and methanolic extracts.

Superoxide radical ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), hydroxyl radical (OH), and singlet oxygen ( $^1O_2$ ) [9–11] are extremely reactive and potentially damaging transient chemical species. Tissue damage, resulting from the imbalance between reactive oxygen species generating and scavenging systems (oxidative stress), has been implicated in the pathology of a number of disorders, such as atherosclerosis, ischemia-reperfusion injury, cancer, malaria, diabetes, inflammatory joint disease, asthma, cardiovascular diseases, cataracts, immune system decline, and could play a role in neurodegenerative diseases and aging processes [10–18].

One of the most common and reliable methods to assay free radical scavenging capacity is the Trolox equivalent antioxidant capacity (TEAC) assay, which was first described by Miller *et al.* (1993) and improved by Re *et al.* [19,20]. It is a decolorization assay in which the pre-formed blue/green radical monocation of 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) ( $ABTS^+$ ) is generated by oxidation of ABTS with potassium persulfate and reduced in the presence of hydrogen-donating antioxidants. TEAC assay values of many compounds and plant extracts have been reported in the literature [21–23]. Since this assay is rather simple, operationally, it can be used in screening studies, especially when  $ABTS^+$  reacts rapidly with antioxidants within 30 min. Moreover,  $ABTS^+$ , which is soluble in both aqueous and organic solvents, is not affected by ionic strength, and is viable over a wide pH range [24]. Due to these favorable qualities it can be used with a wide range of matrices to measure both hydrophilic and lipophilic antioxidant capacities of extracts and body fluids, including typical antioxidants, such as phenolic compounds, and more atypical antioxidants, such as glutathione, uric acid, ascorbic acid,  $\alpha$ -tocopherol,  $\beta$ -carotene, and lycopene [20].

The Folin-Ciocalteu colorimetric assay has been used for many years as a measure of total phenolics in natural products [25]. Its basic mechanism is an oxidation/reduction reaction in which the oxidation of phenols by Folin-Ciocalteu reagent (phosphomolybdic-phosphotungstic acid) yields a colored product with  $\lambda_{max}$  at 765 nm. It is a simple, sensitive, and precise method and can be useful in characterizing and standardizing botanical samples, using gallic acid as reference standard [25].

Many research groups have examined plants as potential sources of natural antioxidants [26–31]. Our research team has capitalized upon access to unique, and largely understudied, plant species from diverse, and at times, extreme habitats and coupled this with the use of two well-established assays for antioxidant activity and total phenolic content, to evaluate the antioxidant activity and total phenolic content of a random sampling of the flora of Jordan. Of the hundreds of previous studies on antioxidant activity of higher plants, few researchers have examined the plants that can be found in the Middle East. In doing so, we have identified a suite of plant species with the antioxidant activity that warrants further investigation, especially for a series of plants wherein the activity could not be attributed to compounds with a phenolic moiety. Moreover, we have established a baseline of antioxidant activities for approximately 3% of the flora of Jordan.

## 2. Results, discussion, and conclusions

### 2.1. Antioxidant activity

The total antioxidant capacity of the aqueous and methanolic extracts of the individual plant species ranged from 20.0 to 916.7  $\mu\text{mol}$  and from 15.1 to 915.6  $\mu\text{mol TE g}^{-1}$  dry weight for aqueous and methanolic extracts, respectively, with little variation between replicates (table 1). Highest levels of antioxidant activity were obtained with the aqueous and methanolic extracts of the Carob tree, *C. siliqua* L. (Fabaceae); namely 916.7 and 915.6  $\mu\text{mol TE g}^{-1}$  dry weight, respectively and from the aqueous extract of the Nettle desert, *Forsskaolea tenacissima* L. (Urtiaceae), namely 813.6  $\mu\text{mol TE g}^{-1}$  dry weight. The high-antioxidant activity of *C. siliqua*, also known as St. John's bread, may be attributed to its high content of polyphenolic compounds [32,33]. However, with respect to *F. tenacissima*, we believe this is the first report of high-antioxidant activity for this plant, which was collected from a unique habitat in the Sudanian biogeographic region in the southern part of Jordan at an elevation of 493 m. This species flourishes mainly in Wadis (Arabic for "valley") of hot deserts, as typified in the lower Jordan Valley and the Dead Sea areas of Jordan [34]. Other plants with high-antioxidant activity were: *Pistacia palaestina* Boiss. (Anacardiaceae), *Viscum cruciatum* Sieb. (Loranthaceae), and *Hypericum triquetrifolium* Turra (Hypericaceae) with values of 549.0/590.9, 460.8/590.5, and 457.4/535.5  $\text{TE g}^{-1}$  dry weight for aqueous/methanolic extracts, respectively.

Based on this collection, approximately 5% of assayed plants showed high levels of antioxidant activity (namely, above 300 and 275  $\mu\text{mol TE g}^{-1}$  dry weight for aqueous and methanolic extracts, respectively), as determined by using *Ginkgo biloba* L. (Ginkgoaceae) as a positive control (311.5 and 275.6  $\mu\text{mol TE g}^{-1}$  dry weight for aqueous and methanolic extracts, respectively). The difference between the antioxidant activity of aqueous and methanolic extracts was statistically significant ( $p < 0.0001$ ).

### 2.2. Total phenolic content

As was the case with antioxidant activity measurements, there was a wide variation in the total phenolic content of the individual plant species, ranging from 4.4 to 78.3 mg and from 2.1 to 52.8 mg gallic acid equivalent (GAE)  $\text{g}^{-1}$  dry weight for the aqueous

Table 1. Antioxidant activity and total phenolic content of aqueous and methanolic extracts of 95 plant species from Jordan.

| Family   | Scientific name   | Antioxidant activity <sup>a</sup><br>( $\mu\text{mol TE g}^{-1}$ dry weight) |                     | Total phenolic content <sup>b</sup><br>( $\text{mg GE g}^{-1}$ dry weight) |                     |             |
|--|---|--|---------------------|--|---------------------|-------------|
|  |   | Aqueous extracts   | Methanolic extracts | Aqueous extracts   | Methanolic extracts |             |
| Acanthaceae                                    | <i>Acanthus syriacus</i> Boiss.                         | 77.0 (0.9%)  | 82.7 (4.0%)         | 15.1 (5.3%)  | 15.4 (0%)           |             |
| Aizoaceae                                      | <i>Aizoon hispanicum</i> L.                             | 48.3 (3.7%)  | 15.1 (2.6%)         | 5.0 (8.0%)   | 2.1 (14.3%)         |             |
|  | <i>Aizoon canariense</i> L.                             | 38.1 (9.9%)  | 15.9 (5.0%)         | 4.4 (22.7%)  | 3.6 (2.7%)          |             |
| Anacardiaceae                                  | <i>Pistacia palaestina</i> (Boiss.)                     | 549.0 (0.5%)   | 590.9 (0.3%)        | 42.5 (5.2%)  | 52.3 (1.9%)         |             |
| Apiaceae                                       | <i>Daucus carota</i> L.                                 | 75.8 (7.0%)  | 49.0 (3.5%)         | 16.9 (4.7%)  | 11.7 (2.6%)         |             |
|  | <i>Ferula communis</i> (L.)                             | 160.7 (1.7%)   | 86.7 (5.4%)         | 24.6 (2.0%)  | 18.4 (2.2%)         |             |
| Asclepiadaceae                                 | <i>Malabaila secacul</i> (Mill) Boiss.                  | 75.6 (0.9%)  | 23.6 (13.1%)        | 13.8 (1.4%)  | 4.0 (7.5%)          |             |
|  | <i>Calotropis procera</i> (Ait)                         | 90.2 (5.3%)  | 50.8 (2.4%)         | 19.4 (3.1%)  | 11.4 (0%)           |             |
| Asteraceae                                     | <i>Gomphocarpus sinaiacus</i> Boiss.                    | 133.3 (1.2%)   | 63.9 (5.3%)         | 25.9 (0.8%)  | 23.8 (7.6%)         |             |
|  | <i>Aaronsohnia factorovskyi</i> Warb & Eig.             | 189.0 (2.3%)   | 214.9 (1.9%)        | 29.1 (2.4%)  | 22.1 (2.7%)         |             |
| Asteraceae                                     | <i>Achillea santolina</i> L.                            | 72.6(0.3%)   | 60.2 (0.3%)         | 20.5 (2.9%)  | 19.2 (1.0%)         |             |
|  | <i>Carthamus tenuis</i> (Boiss & Blanche) Bornm.        | 162.9 (4.1%)   | 61.8 (4.2%)         | 27.8 (4.7%)  | 16.2 (5.6%)         |             |
|  | <i>Centaurea ammocyanus</i> Boiss.                      | 85.0 (7.2%)  | 50.0 (11.0%)        | 18.1 (8.3%)  | 10.9 (22.0%)        |             |
|  | <i>Cichorium pumilum</i> Jacq.                          | 72.7 (3.3%)  | 53.1 (7.3%)         | 16.1 (5.0%)  | 10.6 (0.9%)         |             |
|  | <i>Crepis bulbosa</i> (L.)                              | 55.8 (3.9%)  | 65.0 (0.9%)         | 11.2 (2.7%)  | 11.6 (6.0%)         |             |
|  | <i>Dittrichia viscosa</i> (L.) Greuter                  | 269.8 (1.1%)   | 247.8 (0.9%)        | 57.3 (8.7%)  | 43.9 (3.4%)         |             |
|  | <i>Echinops philistaeus</i> & zohary                    | 100.3 (3.4%)   | 77.8 (4.1%)         | 24.0 (9.2%)  | 19.6 (6.1%)         |             |
|  | <i>Gundelia tournefortii</i> L.                         | 63.6 (8.6%)  | 53.0 (2.8%)         | 16.0 (9.4%)  | 11.0 (7.3%)         |             |
|  | <i>Helichrysum sanguineum</i> (L.) kostel.              | 232.3 (0.5%)   | 164.8 (3.3%)        | 41.3 (9.2%)  | 34.7 (3.2%)         |             |
|  | <i>Iphiona mucronata</i> (Forsk.) Ascherson & Schweinf. | 65.2 (0.5%)  | 46.2 (4.8%)         | 17.3 (2.9%)  | 12.7 (0.8%)         |             |
|  | <i>Matricaria recutita</i> L.                           | 147.6 (0.9%)   | 142.6 (1.0%)        | 20.6 (6.8%)  | 15.1 (3.3%)         |             |
|  | <i>Notobasis syriaca</i> (L.)                           | 64.0 (7.5%)  | 43.8 (5.5%)         | 13.7 (5.8%)  | 8.6 (2.3%)          |             |
|  | <i>Pallenis spinosa</i> (L.) Cass.                      | 50.3 (5.2%)  | 33.8 (0.3%)         | 11.5 (8.7%)  | 7.3 (5.5%)          |             |
|  | <i>Pallenis spinosa</i> (L.) Cass.                      | 55.7 (2.9%)  | 57.1 (6.0%)         | 13.5 (3.7%)  | 10.5 (4.8%)         |             |
|  | <i>Scolymus maculatus</i> L.                            | 70.1 (6.1%)  | 35.8 (1.4%)         | 16.6 (10.8%)   | 9.0 (3.3%)          |             |
|  | <i>Scorzonera syriaca</i> Boiss & Blanche.              | 69.5 (4.0%)  | 44.6 (0.2%)         | 15.1 (9.3%)  | 11.2 (9.8%)         |             |
|  | <i>Varthemia iphionoides</i> Boiss. & BL.               | 176.7 (0.6%)   | 85.3 (1.2%)         | 33.9 (4.7%)  | 20.5 (2.4%)         |             |
|  | <i>Varthemia montana</i> (Vahl) Boiss.                  | 56.9 (0.7%)  | 51.6 (6.2%)         | 10.2 (5.9%)  | 11.1 (0%)           |             |
|  | Boraginaceae  | <i>Alkanna strigosa</i> Boiss. & Hohen.                                      | 157.9 (2.1%)        | 104.2 (5.6%)   | 16.7 (5.4%)         | 11.0 (5.5%) |
|  |   | <i>Anchusa italica</i> Retz.   | 83.3 (1.3%)         | 88.2 (3.3%)  | 12.3 (24.4%)        | 16.2 (0%)   |
|  |   | <i>Anchusa strigosa</i> Banks & Sol.   | 66.7 (2.5%)         | 43.6 (1.4%)  | 10.5 (9.5%)         | 6.1 (1.6%)  |
|  |   | <i>Asperugo procumbens</i> L.  | 189.8 (0.6%)        | 87.5 (2.3%)  | 26.0 (5.0%)         | 15.7 (1.3%) |
|  |   | <i>Echium glomeratum</i> Poir.   | 148.9 (0.9%)        | 108.2 (4.5%)   | 12.8 (19.5%)        | 13.3 (9.0%) |
| <i>Echium judaeum</i> Lacaite                  |   | 70.2 (8.4%)  | 71.3 (0.7%)         | 11.7 (0.9%)  | 11.5 (6.1%)         |             |
| <i>Lappula spinocarpus</i> (Forssk.) O. Kuntze |   | 227.5 (0.5%)   | 124.4 (1.4%)        | 34.0 (1.8%)  | 25.0 (2.4%)         |             |
| <i>Nonea melanocarpa</i> Boiss.                |   | 180.3 (0.8%)   | 129.7 (0.7%)        | 25.1 (6.4%)  | 22.2 (0.9%)         |             |
| <i>Paracaryum rugulosum</i> (D.C) Boiss.       |   | 140.6 (1.0%)   | 98.0 (2.4%)         | 22.8 (6.6%)  | 15.4 (5.8%)         |             |
| <i>Podonosma orientalis</i> (L) Feinbrun.      |   | 176.1 (2.7%)   | 96.6 (3.0%)         | 20.1 (2.5%)  | 19.4 (1.5%)         |             |
| Brassicaceae                                   | <i>Eruca sativa</i> Mill                                | 64.6 (2.5%)  | 34.6 (11.6)         | 19.0 (3.2%)  | 9.5 (4.2%)          |             |
|  | <i>Sinapis alba</i> L.                                  | 49.7 (1.6%)  | 26.5 (1.1%)         | 14.5 (0.7%)  | 8.4 (4.8%)          |             |
|  | <i>Zilla spinosa</i> (L.) Prantl                        | 42.0 (1.0%)  | 23.3 (6.0%)         | 7.3 (6.8%)   | 4.1 (0%)            |             |
| Caesalpiniaceae                                | <i>Cassia italica</i> (Mill)                            | 79.0 (1.5%)  | 53.3 (4.7%)         | 24.0 (0.4%)  | 16.3 (1.2%)         |             |
| Capparaceae                                    | <i>Capparis cartilaginea</i> Decne                      | 91.8 (4.6%)  | 81.1 (4.7%)         | 28.3 (2.8%)  | 20.8 (2.9%)         |             |

(Continued)

Table 1. Continued.

| Family           | Scientific name  | Antioxidant activity <sup>a</sup><br>( $\mu\text{mol TE g}^{-1}$ dry weight) |                     | Total phenolic content <sup>b</sup><br>( $\text{mg GE g}^{-1}$ dry weight) |                     |
|------------------|--|--|---------------------|--|---------------------|
|                  |  | Aqueous extracts   | Methanolic extracts | Aqueous extracts   | Methanolic extracts |
| Chenopodiaceae   | <i>Hammada eigii</i> Iljin                                       | 49.5 (7.9%)  | 30.9 (5.2%)         | 11.3 (6.2%)  | 4.9 (0%)            |
|                  | <i>Salsola jordanicola</i> Eig                                   | 58.9 (1.9%)  | 34.2 (2.0%)         | 9.9 (2.0%)   | 7.2 (0%)            |
|                  | <i>Salsola vermiculata</i> L.                                    | 41.5 (2.7%)  | 27.7 (4.3%)         | 7.0 (1.4%)   | 5.5 (5.5%)          |
|                  | <i>Salsola vermiculata</i> L.                                    | 40.8 (1.2%)  | 32.6 (8.0%)         | 7.8 (3.8%)   | 7.0 (8.6%)          |
| Cistaceae        | <i>Helianthemum lippii</i> L.                                    | 274.2 (2.5%)   | 176.1 (1.8%)        | 30.5 (11.5%)   | 25.0 (6.4%)         |
| Crassulaceae     | <i>Umbilicus intermedius</i> Boiss.                              | 141.8 (4.2%)   | 74.0 (2.7%)         | 18.4 (4.3%)  | 11.9 (0.9%)         |
| Cucurbitaceae    | <i>Bryonia cretica</i> L.  | 120.8 (1.3%)   | 65.0 (6.5%)         | 22.4 (2.7%)  | 13.3 (4.5%)         |
|                  | <i>Bryonia cretica</i> L.  | 170.5 (2.1%)   | 22.0 (10.5%)        | 19.6 (2.6%)  | 5.6 (3.6%)          |
| Dipsacaceae      | <i>Knautia bidens</i> (sm.) Lindley                              | 143.1 (2.4%)   | 78.6 (2.4%)         | 25.7 (5.8%)  | 17.9 (6.1%)         |
| Ephedraceae      | <i>Ephedra alata</i> Decne.                                      | 46.6 (6.9%)  | 60.2 (2.3%)         | 16.2 (2.5%)  | 11.9 (0.8%)         |
| Fabaceae         | <i>Ceratonia siliqua</i> L.                                      | 916.7 (0.2%)   | 915.6 (0.2%)        | 54.2 (3.3%)  | 52.8 (11.4%)        |
|                  | <i>Ononis natrix</i> L.  | 48.0 (5.4%)  | 64.6 (5.6%)         | 9.3 (6.8%)   | 12.3 (1.6%)         |
|                  | <i>Ononis spinosa</i> L. subsp. <i>antiquorum</i> (L.) Arcangeli | 85.9 (4.3%)  | 47.1 (4.9%)         | 18.5 (3.2%)  | 10.6 (15.1%)        |
| Geraniaceae      | <i>Retama raetam</i> (Forssk).                                   | 131.3 (1.8%)   | 67.9 (4.4%)         | 26.4 (18.2%)   | 24.6 (4.1%)         |
|                  | <i>Erodium bryonifolium</i> Boiss                                | 45.3 (2.6%)  | 25.4 (13.0%)        | 15.1 (3.3%)  | 10.8 (3.7%)         |
| Ginkgoaceae      | <i>Ginkgo biloba</i> L. <sup>c</sup>                             | 311.5 (0.4%)   | 275.6 (0.5%)        | 39.0 (9.7%)  | 35.3 (2.8%)         |
| Hypericaceae     | <i>Hypericum triquetrifolium</i> Turra                           | 457.4 (1.7%)   | 535.5 (0.7%)        | 44.6 (4.0%)  | 40.6 (6.4%)         |
| Lamiaceae        | <i>Eremostachys laciniata</i> (L.) Bunge.                        | 85.6 (5.7%)  | 86.8 (7.0%)         | 16.2 (42.6%)   | 19.4 (2.6%)         |
|                  | <i>Marrubium vulgare</i> L.                                      | 76.5 (0.1%)  | 53.9 (9.0%)         | 14.0 (2.9%)  | 11.8 (1.7%)         |
|                  | <i>Phlomis brachyodon</i> (Boiss.) Zohary                        | 59.6 (3.2%)  | 60.4 (3.6%)         | 12.7 (7.9%)  | 15.2 (4.6%)         |
|                  | <i>Phlomis brachyodon</i> Boiss.                                 | 54.2 (0.2%)  | 46.2 (13.9%)        | 9.6 (4.2%)   | 10.9 (23.0%)        |
|                  | <i>Salvia dominica</i> L.  | 29.9 (6.4%)  | 29.4 (5.8%)         | 6.7 (3.0%)   | 6.6 (1.5%)          |
|                  | <i>Salvia hierosolymitana</i> Boiss.                             | 223.9 (0.6%)   | 155.3 (0.8%)        | 33.8 (0.6%)  | 24.4 (0.8%)         |
|                  | <i>Salvia spinosa</i> L. <sup>d</sup>                            | 90.4 (4.8%)  | 74.6 (5.1%)         | 20.8 (6.3%)  | 21.0 (2.4%)         |
|                  | <i>Scutellaria galericulata</i> L.                               | 85.0 (2.2%)  | 77.5 (6.3%)         | 18.8 (11.7%)   | 16.7 (9.0%)         |
|                  | <i>Asparagus aphyllus</i> L.                                     | 65.3 (3.2%)  | 37.8 (5.6%)         | 15.6 (3.8%)  | 9.8 (4.1%)          |
|                  | <i>Scilla hanburyi</i> Baker                                     | 81.0 (2.7%)  | 44.5 (5.2%)         | 17.3 (0.6%)  | 6.9 (7.2%)          |
| Loranthaceae     | <i>Viscum cruciatum</i> Sieb.                                    | 460.8 (0.4%)   | 590.5 (0.2%)        | 39.7 (6.3%)  | 36.5 (4.1%)         |
| Malvaceae        | <i>Alcea acaulis</i> (Cav.) Alef.                                | 20.0 (16.0%)   | 24.2 (1.7%)         | 12.3 (10.6%)   | 5.3 (0%)            |
| Papaveraceae     | <i>Astragalus beershabensis</i> Eig. et Sam.                     | 65.3 (0.9%)  | 29.2 (5.5%)         | 16.3 (1.8%)  | 11.1 (7.2%)         |
|                  | <i>Glaucium aleppicum</i> Boiss. et Hausskn. ex Boiss.           | 161.9 (1.1%)   | 81.2 (3.6%)         | 20.6 (9.2%)  | 18.8 (2.1%)         |
|                  | <i>Lupinus varius</i> L.   | 77.0 (0.6%)  | 69.5 (2.2%)         | 16.8 (0%)  | 13.7 (3.6%)         |
|                  | <i>Roemeria hybrida</i> (L.) DC.                                 | 70.5 (2.6%)  | 58.4 (0.7%)         | 14.9 (22.8%)   | 12.6 (0.8%)         |
| Plantaginaceae   | <i>Plantago lanceolata</i> L.                                    | 29.7 (14.1%)   | 45.3 (8.4%)         | 7.6 (3.9%)   | 9.0 (2.2%)          |
| Polygonaceae     | <i>Rumex pictus</i> Forssk                                       | 161.2 (3.4%)   | 60.6 (6.9%)         | 21.1 (2.8%)  | 10.9 (9.2%)         |
| Resedaceae       | <i>Ochradenus baccatus</i> Del.                                  | 78.2 (3.6%)  | 47.8 (4.0%)         | 20.2 (0.5%)  | 8.8 (2.3%)          |
| Rhamnaceae       | <i>Ziziphus lotus</i> (L.) Lam                                   | 230.2 (0.7%)   | 197.7 (2.3%)        | 30.4 (4.9%)  | 25.0 (11.2%)        |
| Rosaceae         | <i>Amygdalus arabica</i> Oliv.                                   | 234.9 (3.0%)   | 197.6 (0.8%)        | 31.1 (9.0%)  | 31.8 (0.6%)         |
|                  | <i>Amygdalus communis</i> L.                                     | 140.5 (2.3%)   | 179.7 (1.5%)        | 22.7 (2.6%)  | 21.8 (3.2%)         |
| Rutaceae         | <i>Ruta chalepensis</i> L.                                       | 83.6 (6.3%)  | 72.6 (1.5%)         | 21.1 (1.9%)  | 19.5 (3.6%)         |
| Salvadoraceae    | <i>Salvadora persica</i> L.                                      | 38.7 (3.1%)  | 32.0 (1.9%)         | 12.8 (1.6%)  | 10.1 (2.0%)         |
| Scrophulariaceae | <i>Kickxia aegyptiaca</i> (L.) Nabelek                           | 45.7 (3.5%)  | 38.1 (0.8%)         | 16.9 (5.3%)  | 10.7 (1.9%)         |
|                  | <i>Scrophularia xanthoglossa</i> Boiss.                          | 79.3 (4.9%)  | 50.1 (6.4%)         | 18.1 (0.6%)  | 12.3 (9.0%)         |
|                  | <i>Verbascum fruticosum</i> Post                                 | 68.4 (1.6%)  | 46.6 (6.4%)         | 13.4 (1.5%)  | 11.9 (11.8%)        |
|                  | <i>Verbascum jordanicum</i> Murb                                 | 38.0 (4.7%)  | 38.4 (7.6%)         | 9.0 (7.8%)   | 8.7 (5.8%)          |
| Solanaceae       | <i>Hyoscyamus aureus</i> L.                                      | 59.6 (0.7%)  | 52.2 (1.7%)         | 11.9 (2.5%)  | 12.0 (4.2%)         |
| Urticaceae       | <i>Forsskaolea tenacissima</i> L.                                | 813.6 (0.6%)   | 364.3 (1.7%)        | 78.3 (3.1%)  | 39.3 (5.1%)         |
|                  | <i>Urtica pilulifera</i> L.                                      | 49.1 (10.4%)   | 20.8 (3.9%)         | 7.9 (10.1%)  | 4.8 (14.6%)         |

(Continued)

Table 1. Continued.

| Family         | Scientific name  | Antioxidant activity <sup>a</sup><br>( $\mu\text{mol TE g}^{-1}$ dry weight) |                     | Total phenolic content <sup>b</sup><br>(mg GE $\text{g}^{-1}$ dry weight) |                     |
|----------------|--|--|---------------------|---|---------------------|
|                |  | Aqueous extracts   | Methanolic extracts | Aqueous extracts  | Methanolic extracts |
| Zygophyllaceae | <i>Fagonia bruguieri</i> DC.                             | 70.5 (0.9%)  | 33.1 (1.8%)         | 14.4 (2.8%)   | 7.9 (1.3%)          |
|                | <i>Fagonia glutinosa</i><br>Del. Var. grandiflora Boiss. | 66.8 (1.2%)  | 26.1 (1.9%)         | 11.3 (1.8%)   | 7.5 (22.7%)         |
|                | <i>Fagonia mollis</i> Del.                               | 72.5 (0.6%)  | 31.4 (4.5%)         | 24.7 (4.5%)   | 6.7 (0%)            |
|                | <i>Nitraria retusa</i><br>(Forskal) Ascherson            | 44.8 (8.7%)  | 78.0 (4.5%)         | 12.5 (4.8%)   | 15.2 (4.0%)         |
|                | <i>Nitraria retusa</i><br>2(Forskal) Ascherson           | 290.6 (1.4%)   | 200.3 (2.8%)        | 47.8 (3.8%)   | 37.7 (24.9%)        |
|                | <i>Peganum harmala</i> L.                                | 36.2 (10.8%)   | 28.0 (20.4%)        | 8.4 (15.5%)   | 8.0 (25.0%)         |

For both assays, all data are shown as mean (% variation) from two extract replicates, each analyzed in duplicate.

<sup>a</sup>Data are expressed as  $\mu\text{moles}$  of Trolox equivalents per g dry weight.

<sup>b</sup>Data are expressed as mg of Gallic acid equivalents (GAE) per g dry weight.

<sup>c</sup>Positive control for antioxidant activity.

<sup>d</sup>Positive control for total phenolic content.

and methanolic extracts, respectively, with strong reproducibility in the replicate measurements (table 1). Many plant species displayed remarkably high levels of total phenolic content, often with GAE values  $>20 \text{ mg g}^{-1}$  dry weight [35]. This could be compared to *Salvia spinosa*, a genus that is well-known as a rich source of polyphenols with values of 20.8 and 21.0  $\text{mg g}^{-1}$  dry weight for aqueous and methanolic extracts, respectively. The highest values obtained were from the aqueous extracts of *F. tenacissima* and *Dittrichia viscosa* of 78.3 and 57.3  $\text{mg GAE g}^{-1}$  dry weight respectively, and the methanolic extracts of *C. siliqua* and *P. palaestina* of 52.8 and 52.3  $\text{mg GAE g}^{-1}$  dry weight, respectively (table I). Interestingly, neither *D. viscosa* nor *F. tenacissima*, or any other species of the same genera, have been investigated previously for polyphenol content. Several *Pistacia* species are known to be rich in gallotannins and related phenolic compounds [36,37]. With the antioxidant activities, the difference between the aqueous and methanolic extracts regarding total phenolic content was also statistically significant ( $p < 0.0001$ ).

### 2.3. Relationship between antioxidant activity and phenolic content

There was a significant linear correlation between antioxidant activity and total phenolic content for aqueous and methanolic extracts (coefficient  $r = 0.863$  and  $0.850$ , respectively) (figure 1). These results suggested that the phenolic compounds contributed significantly to the antioxidant capacity of the investigated plant species. These results were consistent with the findings of various research groups, who reported positive correlations between total phenolic content and antioxidant activity [26,38].

In addition to the statistical correlation, figure 2 proves to be useful for identification of plants as potential sources for new chemotype antioxidants. A plant with high-antioxidant activity, for which phenolic content *versus* antioxidant activity falls above the regression line (figure 1), is a plant to investigate for novel antioxidants, such as, *C. siliqua* and *V. cruciatum*.




Figure 2. Linear correlation between the total phenolic content and antioxidant activity TEAC. (a) For aqueous extracts: correlation coefficient  $r=0.863$ , coefficient of determination  $r^2=0.744$ . (b) For methanolic extracts: correlation coefficient  $r=0.850$ , coefficient of determination  $r^2=0.723$ .

Moreover, while the total phenolic content of the methanolic extract of *C. siliqua* (52.8 mg GAE g<sup>-1</sup> dry weight) and *P. palaestina* (52.3 mg GAE g<sup>-1</sup> dry weight) is approximately the same, the antioxidant activity of *C. siliqua* (915.6 μmol TE g<sup>-1</sup> dry weight) is 1.5 times higher than that of *P. palaestina* (590.9 μmol TE g<sup>-1</sup> dry weight). We are encouraged by the concept that new, nonphenolic chemotypes with strong antioxidant activity, may be uncovered from those plants where low phenolic content was associated with high-antioxidant activity; such studies are ongoing, in particular, with *V. cruciatum* Sieb., which comes from the high lands in the northern part of Jordan.

The Folin-Ciocalteu assay gives only a crude estimate of the total phenolic compounds in an extract [24]. Hence, this may explain, at least in part, the unequivocal correlation between total phenolic content and antioxidant activity of several plant species (table I). For example, although the total phenolic content of aqueous extract of *D. viscosa* (57.3 mg GAE g<sup>-1</sup> dry weight) was higher than those of *C. siliqua* and *P. palaestina* (54.2 and 42.5 mg GAE g<sup>-1</sup> dry weight), respectively; the corresponding antioxidant activity of *D. viscosa* (269.8 μmol TE g<sup>-1</sup> dry weight) was less than a third with those of *C. siliqua* (916.7 μmol TE g<sup>-1</sup> dry weight) and *P. palaestina* (549.0 μmol TE g<sup>-1</sup> dry weight). Similarly, while the total phenolic content of the methanolic extract of *D. viscosa* (43.9 mg GAE g<sup>-1</sup> dry weight) is higher than that of *V. cruciatum* (36.5 mg GAE g<sup>-1</sup> dry weight), its corresponding antioxidant activity is less than half (247.8 μmol TE g<sup>-1</sup> dry weight in comparison with 590.5 μmol TE g<sup>-1</sup> dry weight). We hope that the ongoing and future research will pursue the identification of the antioxidant compounds from plant species with relatively high-antioxidant activity. A particular focus should be on the plants with high-antioxidant activity and low-phenolic concentrations, as they may prove to yield novel antioxidants. The goal should be to identify and describe new structural classes of natural antioxidant compounds.

### 3. Materials and methods

#### 3.1. General

The total antioxidant capacity assay was performed on a MultiSpec-1501, SHIMADZU® photodiode diode array spectrophotometer (Kyoto, Japan), fitted with Julabo F40, Ultratemp 2000 temperature control. The total phenolic content assay

was carried out using a Spectronic 601 spectrophotometer (Milton Roy Company, USA). Aqueous and methanolic extracts were prepared by using a KARL KOLB water bath shaker (Scientific Technical Supplies, Dreieich, Germany). The incubator B 28 #04-68155 was obtained from Binder GmbH (Bergstr., Tuttlingen, Germany). HPLC grade MeOH and EtOH were obtained from Scharlau Chemie S.A. (Barcelona, Spain) and Fisher Scientific UK limited, (Loughborough, Leicestershire UK), respectively. ABTS<sup>®</sup> was obtained from AppliChem GmbH (Ottoweg, Darmstadt, Germany), and potassium persulfate and Trolox<sup>®</sup> from Acros Organics (New Jersey, USA). Folin-Ciocalteu reagent (2N) was from S.D. Fine-Chem Ltd. (Mumbai, India). Gallic acid monohydrate and sodium carbonate were from Janssen Chemica (Geel, Belgium) and Frutarom Ltd. (Berkhamsted, UK), respectively.

### 3.2. Plant material

The complete range of wild plant samples were collected throughout Jordan, at random, from each of the four biogeographic regions, (figure 1). At each collection point a global positioning system (etrex; Garmin Ltd., Kansas, USA) was used to determine the longitude, latitude, and elevation of the sampling point. These plant samples were identified taxonomically by two of the authors (MS and KAL), and voucher specimens were deposited at the Herbarium Museum of the Faculty of Pharmacy, Jordan University of Science and Technology. The plant raw materials were cleaned of residual soil and air-dried at room temperature. Plants were ground to a fine powder using a laboratory mill, RetschMühle, (RETSCH GmbH, Haan, Germany), passed through a 24 mesh sieve to generate a homogeneous powder, stored at room temperature (22–23°C), and protected from light until required for analyses.

### 3.3. Extraction process

For aqueous extractions, a 250 mg aliquot of each grounded plant material was weighed into a test tube and extracted with 10 mL de-ionized water at 80°C for 1 h in a water bath shaker. After cooling, the extract was centrifuged at  $1507 \times g$  for 10 min at 25°C, and the supernatant was recovered without collection of the pellet and stored at 4°C until used for the TEAC and total phenolic content assays. Methanolic extractions were conducted in a similar manner using 250 mg aliquot of each ground plant material in 10 mL of 80% methanol at 37°C for 3 h in a shaking water bath. After cooling, the extract was centrifuged at  $1507 \times g$  for 10 min, and the supernatant was recovered and stored at 4°C until used for the TEAC and total phenolic content assays. For each plant sample, two replicates of the aqueous and methanolic extracts were prepared [26].

### 3.4. Measurement of totalantioxidant activity

The antioxidant capacity assay was carried out using the improved ABTS<sup>+</sup> method as described by Re *et al.* [20], using Trolox as a standard. Total antioxidant activity was expressed in terms of Trolox equivalent antioxidant capacity (TEAC,  $\mu\text{mol}$  Trolox equivalents per g dry weight of plant) [20].

### 3.5. Measurement of total phenolic content

Total phenolic content was estimated using the Folin-Ciocalteu colorimetric method based on the procedure of Singleton and Rossi, using gallic acid as a standard [25]. Briefly, 50  $\mu\text{L}$  aliquots from each of the replicates were mixed with 450  $\mu\text{L}$  of DI  $\text{H}_2\text{O}$  and 2.5 mL of 0.2 N Folin-Ciocalteu reagent. After 5 min, 2 mL of saturated sodium carbonate ( $75 \text{ g L}^{-1}$ ) was added. The absorbance of the resulting blue solution was measured at 765 nm after incubation at  $30^\circ\text{C}$  for 1.5 h with intermittent shaking. Quantitative measurements were performed based on a six point standard calibration curve of 20, 100, 200, 300, 400, 500  $\text{mg L}^{-1}$  of gallic acid in 80% methanol. The total phenolic content is expressed as gallic acid equivalents (GAE) in milligrams per gram dry material.

### 3.6. Statistical analysis

For both assays, the entire data in table 1 is shown as mean (% variation) from two extract replicates, each analyzed in duplicate. Correlation and regression analysis of antioxidant activity ( $Y$ ) versus the total phenolic content ( $X$ ) was carried out by using Microsoft Office Excel 2003, and the quality control was checked and matched with correlation and regression analyses in SAS 9.1 by using the linear regression command, proc reg. Paired  $t$ -test was applied to test for significant differences between aqueous and methanolic extracts for antioxidant activity and total phenolic content.

### 3.7. Geographic information systems

The data for figure 1 was created and generated by importing the latitude/longitude locations of plant sample sites and nature preserves from the GPS device into a GIS layer by using ArcGIS 9.1 technology (ESRI; Redlands, CA). Identifiers were assigned to each point and the attributes describing each location were assigned from field data sheets. The GIS layer of biogeographical regions was derived from the work of Al-Eisawi [1]. The standard GIS sources for country administrative boundaries, major hydrography, and major cities were used for the remaining map content.

## Acknowledgements

This work was supported in part by the Fogarty International Center of the National Institutes of Health with co-funding from the National Institute of Allergy and Infectious Diseases, the National Institute of Drug Abuse, and the National Science Foundation under grant number R21 TW006628 for the International Cooperative Biodiversity Groups. The authors from the Jordan University of Science and Technology acknowledge the kind financial support of the Deanship of Scientific Research. We would also like to thank Jeanne M. Game for assistance in creating Map 1.

## References

- [1] D.M. Al-Eisawi. *Field Guide to Wild Flowers of Jordan and Neighbouring Countries*, pp. 1–12, Jordan Press, Foundation Al-Rai, Amman, Jordan (1998).
- [2] N. Feinbrun-Dothan. *Flora Palestina*, p. 43, The Israel Academy of Sciences and Humanities, Jerusalem (1986).
- [3] B. Abu-Irmaileh, F.U. Afifi. *Dirasat, Med. Biol. Sci.*, **27**, 53 (2000).
- [4] S. Al-Khalil. *Int. J. Pharmacogn.*, **33**, 317 (1995).
- [5] Anonymous. Conservation and Sustainable Use of Biological Diversity in Jordan, The Hashemite Kingdom of Jordan, Amman (2001).
- [6] F. Alali, A.S. Ma'aya'h, A. Alkofahi, A. Qandil, C. Li, J.P. Burgess, M.C. Wani, N.H. Oberlies. *Nat. Prod. Commun.*, **1**, 95 (2006).
- [7] F.Q. Alali, T. El-Elimat, C. Li, A. Qandil, A. Alkofahi, K.K. Tawaha, J.P. Burgess, Y. Nakanishi, D.J. Kroll, H.A. Navarro, J.O.F. III, M.C. Wani, N.H. Oberlies. *J. Nat. Prod.*, **68**, 173 (2005).
- [8] F.Q. Alali, K. Tawaha, T. El-Elimat, R. Qasaymeh, C. Li, J. Burgess, Y. Nakanishi, D.J. Kroll, M.C. Wani, N.H. Oberlies. *Nat. Prod. Res.*, **20**, 558 (2006).
- [9] O.I. Aruoma. *Food Chem. Toxicol.*, **32**, 67183 (1994).
- [10] M. Martínez-Cayuela. *Biochimie*, **77**, 147 (1995).
- [11] C. Schöneich. *Exp. Gerontol.*, **34**, 19 (1999).
- [12] T.M. Florence. *Aust. New Zealand J. of Ophthalmol.*, **23**, 3 (1995).
- [13] D. Harman. *Age*, **16**, 23 (1993).
- [14] P.C.H. Hollman, M.G.L. Hertog, M.B. Katan. *Food Chem.*, **57**, 43 (1996).
- [15] A.S. Meyer, M. Heinonen, E.N. Frankel. *Food Chem.*, **61**, 71 (1998).
- [16] T. Nakagami, N. Nanaumi-Tamura, K. Toyomura, T. Nakamura, T. Shigehisa. *J. Food Sci.*, **60**, 653 (1995).
- [17] W.S. Waring. *Proc. R. Coll. Physicians Edinb.*, **31**, 288 (2001).
- [18] I.S. Young, J.V. Woodside. *J. Clinical Pathol.*, **54**, 176 (2001).
- [19] N.J. Miller, C. Rice-Evans, M.J. Davies, V. Gopinathan, A. Milner. *Clinical Sci.*, **84**, 407 (1993).
- [20] R. Re, N. Pellegrini, A. Proteggente, A. Pannala, M. Yango, C. Rice-Evans. *Free Radical Biol. Med.*, **26**, 1231 (1999).
- [21] I.L. Nielsen, G.R. Haren, E.L. Magnussen, L.O. Dragsted, S.E. Rasmussen. *J. Agri. Food Chem.*, **51**, 5861 (2003).
- [22] P. Pellergrini, M. Serafini, B. Colombi, D.D. Rio, S. Salvatore, M. Bianchi, F. Brighenti. *J. Nutr.*, **133**, 2812 (2003).
- [23] A.R. Proteggente, A.S. Pannala, G. Paganga, L.V. Buren, E. Wagner, S. Wiseman, F.V.D. Put, C. Dacombe, C.A. Rice-Evans. *Free Radical Res.*, **36**, 21733 (2002).
- [24] R.L. Prior, X. Wu, K. Schaich. *J. Agri. Food Chem.*, **53**, 4290 (2005).
- [25] V.L. Singleton, J.A. Rossi. *Am. J. Enol. Viticulture*, **16**, 144 (1965).
- [26] Y. Cai, Q. Luo, M. Sun, H. Corke. *Life Sci.*, **74**, 2157 (2004).
- [27] V. Katalinic, M. Milos, T. Kulisic, M. Jukic. *Food Chem*, **94**, 5507 (2006).
- [28] M. Oktay, İ. Gülçin, Ö.İ. Küfrevioğlu. *Lebensmittel-Wissenschaft Und-Technologie-Food Sci. Tech.*, **36**, 26371 (2003).
- [29] S.P. Wong, L.P. Leong, J.H.W. Koh. *Food Chem.*, **99**, 77583 (2006).
- [30] E. Middleton, C. Kandaswami. *In The Flavonoids: Advances in Research since 1986*, pp. 619–20, Chapman and Hall, London, UK (1994).
- [31] G.E. Trease, W.C. Evans. *Trease and Evans' Pharmacognosy*, 13th Edn, pp. 378–423, Bailliere Tindall, London and Philadelphia (1989).
- [32] S. Marakis. *J. Food Sci. Technol.*, **33**, 365 (1996).
- [33] R.W. Owen, R. Haubner, W.E. Hull, G. Erben, B. Spiegelhalter, H. Bartsch, B. Haber. *Food Chem. Toxicol.*, **41**, 1727 (2003).
- [34] N. Feinbrun-Dothan. *Flora Palestina*, pp. Maps 1 and 2, The Israel Academy of Sciences and Humanities, Jerusalem (1986).
- [35] M.P. Kähkönen, A.I. Hopia, H.J. Vuorela, J.P. Rauha, K. Pihlaja, T.S. Kujala, M. Heinonen. *J. Agri. Food Chem.*, **47**, 3954 (1999).
- [36] M.C. Barotto, M. Tattini, C. Galardi, P. Pinelli, A. Romani, F. Visioli, R. Basosi, R. Pogni. *Free Radical Res.*, **37**, 405 (2003).
- [37] A.J. Hou, L.Y. Peng, Y.Z. Liu, Z.W. Lin, H.D. Sun. *Planta Medica*, **66**, 6246 (2000).
- [38] W. Zheng, S.Y. Wang. *J. Agri. Food Chem.*, **49**, 5165 (2001).